


Osteoarthritis in Cats

An Overview of
Causes, Diagnostics,
and Therapy



Osteoarthritis in Cats


An Overview of Causes, Diagnostics, and Therapy



Godshorn (Germany)

HEALTHY JOINT

ARTHRITIC JOINT



ERODED
CARTILAGE

BONE
SPURS

INFLAMED
SYNOVIUM

Chapter 1

The History of Awareness

From Age-Related Ailment

to Chronic Pain Disorder

1.1 Historical Misconception and Paradigm Shift

For many years, osteoarthritis in cats was viewed as nothing more than a harmless sign of aging. It was assumed that joints simply became stiffer over time and that this was part of the natural life cycle. The possibility that cats might actually suffer was rarely considered. A major reason lies in feline behavior: cats seldom show pain openly because, in the wild, displaying weakness can be dangerous. As a result, many cats appear outwardly normal even when they are already experiencing significant discomfort.

Diagnostic challenges added to the problem. Radiographs often revealed joint changes, but these did not always match the cat's clinical presentation. Some cats with severe radiographic findings seemed stable, while others with only mild visible changes showed clear signs of pain. This inconsistency contributed to osteoarthritis being underestimated for decades.

A turning point came with newer studies. Beginning in the 2010s, research showed that a large percentage of cats over the age of twelve exhibit signs of osteoarthritis — roughly 90 percent.

This made the true extent of the disease unmistakable and highlighted how much suffering had previously gone unnoticed. Today, even subtle behavioral changes are taken seriously. Reduced jumping ability, lower activity levels, a poorly groomed coat, or relocating a sleeping spot to an easier-to-reach area are now recognized as early warning signs.

1.2 Pathophysiology and Risk Factors

Osteoarthritis develops through the gradual breakdown of articular cartilage. This cartilage acts as a cushion between bones. As it becomes thinner or uneven, friction increases, triggering inflammation within the joint. This inflammation causes pain and promotes further structural damage. Although the existing damage cannot be reversed, the progression of the disease can be slowed.

In cats, primary osteoarthritis is most common and closely associated with aging. Modern domestic cats live significantly longer than in the past, which increases the overall prevalence. Yet early changes can begin in much younger animals. Osteoarthritis is therefore typically a long-term process that evolves over many years.

Secondary osteoarthritis occurs when a joint has been previously damaged — for example through injury, developmental abnormalities, or chronic inflammation. Another major risk factor is excess weight. Obesity puts additional strain on the joints and amplifies inflammatory processes throughout the body.

Key risk factors include:

- advanced age
- overweight and low muscle mass
- previous injuries
- anatomical misalignments or developmental disorders
- recurring or chronic inflammation
- possible genetic influences



Chapter 2

The Scientific Basis of Pain

From Inflammation to

Neural Sensitization

2.1 Mechanisms of Chronic Pain Development

Chronic joint pain in cats arises from several interconnected processes. As articular cartilage breaks down, friction within the joint increases. This friction irritates the joint capsule and synovial membrane, leading to inflammation that causes swelling, warmth, and pressure sensitivity.

Over time, the biochemical environment within the joint also changes. A common issue is sustained tissue acidosis, which raises the sensitivity of pain receptors and intensifies local pain. At the same time, nerve endings in the joint become more easily excitable and react more strongly to movement or mechanical load.

If these processes persist, central sensitization can develop. In this state, the nervous system becomes increasingly responsive, and stimuli that would normally be mild may trigger significant pain. Some cats even react sensitively to light touch or begin to withdraw socially. This mechanism explains why osteoarthritic pain can appear more severe than structural joint changes alone would suggest.

2.2 The Nerve Growth Factor (NGF) as a Key Molecule

A major scientific breakthrough was the discovery of the role of Nerve Growth Factor (NGF). This molecule is released in higher amounts within inflamed or damaged joints. It binds to specific receptors on nociceptive nerve fibers and increases their sensitivity, resulting in more frequent and more intense pain signaling.

NGF is involved not only in transmitting pain but also in promoting hypersensitivity. Traditional pain medications were not able to adequately target this particular component of chronic pain. Only a detailed understanding of the NGF signaling pathway made it possible to develop modern, targeted therapies that directly block this mechanism.

Key characteristics of NGF-mediated pain:

- heightened sensitivity of nociceptive fibers
- increased tendency of the nervous system to generate pain signals
- significant contribution to the development of chronic pain
- limited effectiveness of many traditional anti-inflammatory drugs against this mechanism



Chapter 3

The Evolution of Diagnostics

Detecting Hidden Pain

3.1 From Subjective Impressions to Objective Assessment

Diagnosing osteoarthritis pain in cats has long been a challenge. Cats rarely show pain clearly, especially not in unfamiliar environments. In the veterinary clinic, many cats appear calm, tense, or simply stressed, which makes orthopedic assessment difficult. As a result, pain was often overlooked for decades.

Traditionally, diagnosis relied on physical examination, radiographs, and the owner's general impressions. This approach was unreliable. Radiographic images reveal structural changes, but these do not always correlate with the cat's actual level of discomfort. Some cats experience significant pain without obvious radiographic abnormalities, while others with marked changes show few outward symptoms. This mismatch contributed to the persistent underestimation of osteoarthritis pain.

Over time, it became evident that everyday behavior at home provides far more accurate information than the cat's presentation in the clinic. Stress behaviors in the exam room can mask or overshadow signs of orthopedic disease.

Modern diagnostics therefore focus much more heavily on observations from the home environment.

3.2 The Feline Musculoskeletal Pain Index (FMPI)

A major advancement was the development of the Feline Musculoskeletal Pain Index (FMPI) by North Carolina State University. The FMPI is a validated questionnaire specifically designed to assess chronic joint pain in cats and is currently considered the most important standardized tool in this field.

The questionnaire covers typical daily activities such as jumping, climbing stairs, grooming, playing, and overall activity. Owners rate how well the cat performs each of these tasks. This creates an accurate picture of the cat's true mobility and functional behavior. The FMPI effectively bridges the gap between clinical observation and everyday life.

A major benefit is the elimination of stress-related distortions. At home, cats behave naturally, making these assessments far more reliable.


Typical aspects evaluated in the FMPI include:

- ability to jump, especially to higher surfaces
- stair climbing and general mobility
- extent and quality of grooming
- play behavior and activity level
- interactions with familiar people


The FMPI not only supports initial diagnosis but is also valuable for monitoring treatment progress over time.



Nerve
Receptor



Monoclonal
Antibody



Nerve
Receptor

Chapter 4

The History of Therapy

Challenges and Breakthroughs

4.1 The Era of Pharmacologic Restriction

For many years, options for treating chronic pain in cats were extremely limited. This was largely due to the unique metabolic physiology of the feline body. Cats lack a major detoxification pathway — glucuronidation — which allows many drugs to be safely metabolized in humans and dogs. Without this pathway, certain medications remain in the system much longer and can cause toxic effects.

As a result, many well-known human pain relievers are dangerous for cats. Acetaminophen is the most prominent example: even very small doses can be life-threatening and lead to severe liver damage. Many common NSAIDs routinely used in human medicine are also unsuitable for feline patients. For decades, these risks led to extreme caution in veterinary pain management. Out of fear of side effects, cats were often undertreated and left without adequate pain control.

4.2 Establishment of Modern NSAID Safety Concepts

As understanding of osteoarthritis and its consequences grew, it became clear that untreated chronic pain poses a serious health risk on its own.

Modern NSAIDs formulated specifically for cats have therefore become an essential part of therapy — but only with strict safety principles in place.

Key principles for safe NSAID use in cats:

- use only veterinary-approved products
- monitor kidney and liver values regularly
- employ the lowest effective dose
- adjust treatment individually based on the cat's condition
- exercise caution with comorbidities such as CKD or liver disease
- avoid human NSAIDs entirely

Following these guidelines allows controlled, effective, and long-term pain management. For many cats, this results in significantly improved mobility and overall quality of life.



Chapter 5

The Present – Cellular and Biological Advances in Pain Management

5.1 Monoclonal Antibodies – Blocking the NGF Pathway

The most significant breakthrough of recent years is the introduction of monoclonal antibodies for managing chronic osteoarthritic pain. These biologic agents target very specific components of the pain pathway. They inhibit Nerve Growth Factor (NGF), a key molecule involved in generating and amplifying chronic pain.

For cats, the first approved anti-NGF antibody is frunevetmab (brand name Solensia). It binds to NGF within affected joint tissues, preventing it from attaching to receptors on pain-transmitting nerve fibers. As a result, nerve hypersensitivity decreases, and pain is relieved in a sustained and targeted manner.

A major advantage of this therapy is its excellent safety profile. Monoclonal antibodies are not metabolized through the liver but are broken down naturally as proteins. This means they place minimal strain on the cat's detoxification pathways. They are therefore well suited for older cats or those with underlying organ disease.

The once-monthly injection also improves compliance, eliminating the need for daily medication.

5.2 Adjunctive Medications for Pain Processing Control

In addition to monoclonal antibodies, other medications are used to address specific aspects of altered pain processing. These are typically added when osteoarthritis is complex or when pain is already well established.

Commonly used adjunctive medications include:

– Amantadine:

An NMDA receptor antagonist that can reduce central sensitization. It is especially useful when pain persists despite primary therapy.

– Gabapentin:

Often used for neuropathic pain. It decreases nerve excitability and can provide meaningful relief in difficult cases.

Neither of these medications is a classic anti-inflammatory. Instead, they target the changed neural pain pathways within the central nervous system. Their benefits may take several days to develop, but they are valuable additions in more severe or refractory cases.

These modern pharmacologic approaches have fundamentally changed osteoarthritis treatment. For the first time, a therapeutic principle is available that not only dampens symptoms but interrupts the mechanism of chronic pain at its source.

MULTIMODAL CONCEPT



WEIGHT





NUTRITION



MEDICATION



ENVIRONMENT



Chapter 6

Multimodal Management

Integrating Environment,

Nutrition, and Rehabilitation

6.1 Weight Management and Nutrition

One of the most important non-pharmacologic measures in osteoarthritis care is weight management. Excess weight places additional stress on the joints and intensifies inflammatory processes throughout the body. Even moderate weight loss can significantly reduce pain and improve mobility. A controlled, calorie-adjusted feeding plan is therefore a central component of treatment.

Targeted nutritional support also plays a role. Omega-3 fatty acids — especially EPA and DHA — have anti-inflammatory properties and can complement the effects of other pain therapies. Chondroprotective supplements such as green-lipped mussel powder or specific glycosaminoglycans support joint function. While they do not halt disease progression, they may improve joint resilience and reduce discomfort.

Veterinary-guided nutritional counseling is especially valuable when a cat is overweight or has additional medical conditions.

6.2 Environmental Modifications and Physical Therapy

Adapting the home environment can make daily life far easier for a cat with osteoarthritis. Many helpful measures are simple to implement and offer immediate benefits. Ramps or steps can assist the cat in reaching favorite resting places, and beds with low entry points improve accessibility. Litter boxes with shallow sides are much more comfortable for cats with joint pain.

Soft, well-padded resting areas help reduce pressure on sensitive joints. At the same time, the living space should be arranged so that the cat can access key locations without needing to jump or exert excessive effort.

Physical therapy can provide additional support. Gentle massage, warm or cool compresses, and carefully guided mobility exercises can improve blood flow, relax muscles, and reduce pain. Individualization is crucial — treatments must be comfortable for the cat and must never cause strain or additional discomfort.

The success of osteoarthritis therapy depends not on a single intervention but on a combination of strategies. A multimodal approach takes the entire well-being of the cat into account, resulting in substantially improved quality of life.



Chapter 7

Complementary Therapies

and Alternative Approaches

7.1 Supportive Physical Techniques

Beyond medication, several physical modalities can help make life more comfortable for cats with osteoarthritis. These methods do not replace pain therapy, but they can provide meaningful support and symptom relief.

Heat therapy:

Heat relaxes tense muscles, improves circulation, and can ease stiff joints. Short, mild applications — such as a lukewarm hot water bottle wrapped in a towel — are effective. It is essential to avoid overheating and apply heat only indirectly.

Cold therapy:

Cold can help reduce swelling and pain during acute flare-ups. As with heat, cold should be used briefly, gently, and never applied directly to the skin.

Massage and gentle mobilization:

Light massages can help relax the muscles and improve mobility. They must be done carefully, as many cats are sensitive to excessive handling. Short, calm movements are sufficient. Overstretching must be strictly avoided.

7.2 Laser, Electrical, and Magnetic Therapies

Some veterinary practices offer additional technical modalities that may support pain management.

Low-level laser therapy:

The “cold laser” can help reduce inflammation and stimulate tissue healing. Many cats tolerate this treatment well. The effect varies individually but is often perceived as soothing.

TENS (transcutaneous electrical nerve stimulation):

This method uses mild electrical impulses to influence pain signaling. It requires experience and proper instruction. Not all cats are suitable candidates, but some benefit from reduced pain sensitivity.

Magnetic field therapy:

Used in some practices as a complementary option. The scientific evidence is limited, but some cats respond positively to regular sessions. It is not sufficient as a standalone treatment but may be used as an adjunct.

7.3 Acupuncture and Gold Implantation

Acupuncture:

One of the most well-known alternative approaches. Some cats respond well to gentle needle stimulation, particularly when muscle tension contributes to their discomfort. Responses vary, and treatment must be performed by trained professionals.



Gold implantation:

Small pieces of gold are placed near affected joints. The method is debated because scientific evidence is limited. Some cats improve, while others show little change. Thorough veterinary evaluation is essential before considering the procedure.

7.4 Supplements and Natural Therapies

Many owners seek natural options to support treatment. Some can be helpful, while others are ineffective or even harmful.

Nutritional supplements with potential benefits:

- Omega-3 fatty acids (EPA, DHA)
- green-lipped mussel
- select joint-support complexes (veterinary monitored)
- antioxidants included in specialized diets

These supplements may help moderate inflammation or support cartilage health. They do not replace medical pain therapy and work best as part of an overall plan.

Herbal remedies:

Some plants have anti-inflammatory properties, but many herbs are toxic to cats. Any use must be discussed with a veterinarian.

Homeopathy, Bach flowers, and similar methods:

Scientific evidence is minimal. Some owners report subjective improvements, mostly in emotional well-being. These methods may be used as long as they do not replace medical treatment.

7.5 Key Principle

All complementary therapies can offer support, but none should replace medically necessary treatment.

Osteoarthritis is a chronic, progressive disease that requires a structured, evidence-based plan. Alternative approaches are most effective when used thoughtfully, in proper doses, and with consistent veterinary oversight.



Chapter 8

Conclusion and Outlook

Osteoarthritis in cats was overlooked for a long time. Many animals suffered silently because their pain was either not recognized or dismissed as a normal part of aging. Today, even subtle behavioral changes are taken seriously. New diagnostic tools such as the FMPI make it easier to identify and monitor chronic pain in everyday life.

Therapeutic options have also advanced significantly. Modern pain medications — especially monoclonal antibodies — enable targeted treatment of central pain mechanisms. Adjunctive medications like amantadine or gabapentin help when the nervous system is particularly sensitized.

At the same time, classic measures remain essential: controlled weight management, joint-supportive nutrition, environmental adjustments, and physiotherapeutic support are fundamental components of comprehensive care.

Future progress will likely focus on even earlier detection as well as new biological agents that further refine pain modulation. The goal is to slow disease progression and provide cats with osteoarthritis the best possible comfort and mobility throughout their lives.

Osteoarthritis cannot be cured, but it can be effectively managed. With proper knowledge, careful observation, and a well-coordinated treatment plan, veterinarians and caregivers can work together to meaningfully improve the long-term well-being of affected cats.



For your notes

For your notes



Cat Man (Booklet's)